



TITLE:

Keller-Segel System and the Concentration Lemma(Variational Problems and Related Topics)

AUTHOR(S):

NAGAI, Toshitaka; SENBA, Takasi; SUZUKI, Takashi

CITATION:

NAGAI, Toshitaka ...[et al]. Keller-Segel System and the Concentration Lemma(Variational Problems and Related Topics). 数理解析研究所講究録 1998, 1025: 75-80

ISSUE DATE:

1998-02

URL:

<http://hdl.handle.net/2433/61743>

RIGHT:

Keller-Segel System and the Concentration Lemma

Department of Mathematics, Kyushu Institute of Technology

Toshitaka NAGAI
(永井 敏隆)

Department of Applied Mathematics, Miyazaki University

Takasi SENBA
(仙葉 隆)

Department of Mathematics, Osaka University

Takashi SUZUKI
(鈴木 貴)

1 Introduction

We consider time-global existence and blow-up of solutions of the following system related to chemotaxis

$$(P) \quad \begin{cases} u_t = \nabla \cdot (\nabla u - \chi u \nabla v), & x \in \Omega, t > 0, \\ 0 = \Delta v - \gamma v + \alpha u, & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, & x \in \partial\Omega, t > 0, \\ u(\cdot, 0) = u_0, & x \in \Omega. \end{cases}$$

Here Ω is a bounded domain in \mathbf{R}^2 with smooth boundary $\partial\Omega$, χ , γ and α are positive constants and u_0 is a non-negative smooth function on $\bar{\Omega}$.

There exists a unique solution (u, v) to (P) defined on a maximal interval of existence $[0, T_{max})$, which is smooth in $x \in \bar{\Omega}$ and $0 < t < T_{max}$. If $u_0 \not\equiv 0$ in Ω , the solution satisfies that $u(x, t) > 0$, $v(x, t) > 0$ for $(x, t) \in \Omega \times (0, T_{max})$. If $T_{max} < \infty$, we can observe the following.

Proposition 1 *If $T_{max} < \infty$, then the following relations hold.*

- (i) $\lim_{t \rightarrow T_{max}} \|u \log u\|_{L^1(\Omega)} = \infty$
- (ii) $\lim_{t \rightarrow T_{max}} \|\nabla v\|_{L^2(\Omega)} = \infty$.
- (iii) For $a > \chi/2$, then $\lim_{t \rightarrow T_{max}} \int_{\Omega} e^{av(x,t)} dx = \infty$.

Then, if $T_{max} < \infty$, we have that

$$\lim_{t \rightarrow T_{max}} \|u(\cdot, t)\|_{L^\infty(\Omega)} = \lim_{t \rightarrow T_{max}} \|v(\cdot, t)\|_{L^\infty(\Omega)} = \infty,$$

which we mean that the solution blows up in finite time.

Let L be an arbitrary positive constant and let $D_L = \{x \in \mathbf{R}^2 | |x| < L\}$. We have the following results.

Theorem 1 *Suppose*

$$\Omega = D_L \quad \text{and} \quad \|u_0\|_{L^1(D_L)} < 8\pi/(\alpha\chi). \quad (1)$$

Let $u_0(x) = u_0(-x)$ on D_L . Then (P) admits a unique classical solution (u, v) on $\overline{D_L} \times (0, \infty)$ satisfying

$$\sup_{t \geq 0} \{ \|u(\cdot, t)\|_{L^\infty(D_L)} + \|v(\cdot, t)\|_{L^\infty(D_L)} \} < \infty.$$

Definition 1 We say that q is a blow-up point of u if there exists $\{t_k\}_{k=1}^\infty \subset [0, T_{\max})$ and $\{x_k\}_{k=1}^\infty \subset \overline{\Omega}$ satisfying $u(x_k, t_k) \rightarrow \infty$, $t_k \rightarrow T_{\max} < \infty$ and $x_k \rightarrow q \in \overline{\Omega}$ as $k \rightarrow \infty$. We denote the set of all blow-up points of u by \mathcal{B} .

Theorem 2 Let (1) hold. Let a_* be a root of $a_* - \chi/2 - \|u_0\|_{L^1(\Omega)} \alpha a_*^2 / 16\pi = 0$ such that $a_* < \chi$. If $T_{\max} < \infty$, then there exists a point $q \in \mathcal{B} \cap \partial\Omega$ satisfying

$$\limsup_{t \rightarrow T_{\max}} \int_{\Omega \cap B(q, \varepsilon)} u(x, t) dx \geq \frac{2\pi}{a_* \alpha} \quad \text{for any } \varepsilon > 0.$$

Definition 2 For $q \in \mathcal{B}$, we say that q is an isolated blow-up point if there exists $\delta > 0$ such that

$$\sup\{u(x, t) \mid 0 \leq t < T_{\max} \text{ and } x \in \overline{B(q, \delta) \setminus B(q, \varepsilon)} \cap \Omega\} < \infty \quad \text{for any } \varepsilon \in (0, \delta).$$

We denote the set of all isolated blow-up points of u by \mathcal{B}_I .

Theorem 3 Suppose $T_{\max} < \infty$. Then the following properties hold.

For $q \in \mathcal{B}_I$, there exist two positive constants δ , $m \geq m_*$ and a non-negative function $f \in L^1(E(q, \delta)) \cap C(E(q, \delta) \setminus \{q\})$ such that

$$w^* - \lim_{t \rightarrow T_{\max}} u(\cdot, t) = m\delta_q + f \quad \text{in } \mathcal{M}(E(q, \delta)),$$

where $E(q, \delta) = \overline{B(q, \delta)} \cap \Omega$,

$$m_* = \begin{cases} 4\pi/(\alpha\chi) & \text{if } q \in \partial\Omega, \\ 8\pi/(\alpha\chi) & \text{if } q \in \Omega \end{cases}$$

and $\mathcal{M}(S)$ is the Banach space consisting of all Radon measures on a compact Hausdorff space S with the usual norm.

For a set K , we denote the number of elements of K by $\#K$. The following corollary is an immediate consequence of Theorem 3.

Corollary 1 Suppose $T_{\max} < \infty$. Then \mathcal{B}_I satisfies that

$$\#\{\mathcal{B}_I \cap \Omega\} + \frac{1}{2} \#\{\mathcal{B}_I \cap \partial\Omega\} \leq \frac{\alpha\chi}{8\pi} \|u_0\|_{L^1(\Omega)}.$$

The following corollary is an immediate consequence of Theorem 3 and [6].

Corollary 2 Suppose $\Omega = D_L$ and that u_0 be radially symmetric. If $\int_{D_L} u_0(x) dx > 8\pi/(\alpha\chi)$ and $\int_{D_L} u_0(x) |x|^2 dx$ is sufficiently small, then $T_{\max} < \infty$ and there exist a positive constant $m \geq 8\pi/(\alpha\chi)$ and a non-negative function $f \in L^1(D_L) \cap C(\overline{D_L} \setminus \{0\})$ such that

$$w^* - \lim_{t \rightarrow T_{\max}} u(\cdot, t) = m\delta_0 + f \quad \text{in } \mathcal{M}(\overline{D_L}).$$

2 Proof of Theorem 1

Lemma 1 *Let (u, v) be a solution to (P). Put*

$$W(t) = \int_{\Omega} \left\{ u \log u - \frac{\chi}{2\alpha} (|\nabla v|^2 + \gamma v^2) \right\} dx.$$

Then, it follows that

$$\frac{d}{dt} W(t) + \int_{\Omega} u |\nabla (\log u - \chi v)|^2 dx = 0 \quad \text{for } t \in (0, T_{max}).$$

Proof of Lemma 1: Multiplying $\log u - \chi v$ by the first equation of (P) and using the second equation of (P), then we have this lemma.

Lemma 2 *Suppose that (u, v) is a solution to (P). Let a be an arbitrary positive constant and let $M = \|u_0\|_{L^1}$. Then, the inequality*

$$a \int_{\Omega} uv dx \leq \int_{\Omega} u \log u dx + M \log \left(\int_{\Omega} e^{av} dx \right) - M \log M$$

holds for $0 \leq t < T_{max}$.

Proof of Lemma 2. Let

$$\mu = \int_{\Omega} e^{av} dx \text{ and } \psi = \frac{M}{\mu} e^{av}.$$

Then, we have that

$$\begin{aligned} 0 &= -\log \left(\int_{\Omega} \frac{\psi u}{u \mu} dx \right) \\ &\leq \int_{\Omega} \left(-\log \frac{\psi}{u} \right) \frac{u}{M} dx, \end{aligned}$$

by which together with Jensen's inequality we get this lemma.

Proposition 2 *If w is a function on $\overline{D_L}$ satisfies that $w \in C^1(D_L)$, $w(x) = w(-x)$ on ∂D_L and*

$$\frac{\partial w}{\partial n} = 0 \quad \text{on } \partial D_L,$$

then there exists absolute constants C and K such that

$$\log \left(\int_{\Omega} e^w dx \right) \leq \frac{1}{16\pi} \int_{\Omega} |\nabla w|^2 dx + C \|w\|_{L^1} + K.$$

Proof of Theorem 1. By Lemmas 1, 2 and the second equation of (P), we have that

$$\left(a - \frac{\chi}{2}\right) \frac{1}{\alpha} \int_{\Omega} (|\nabla v(x, t)|^2 + \gamma |v(x, t)|^2) dx \leq W(0) + M \log \left(\int_{\Omega} e^{av(x, t)} dx \right) - M \log M,$$

by which together with Proposition 2 it follows that

$$\left\{ \left(a - \frac{\chi}{2}\right) \frac{1}{\alpha} - \frac{Ma^2}{16\pi} \right\} \int_{\Omega} (|\nabla v(x, t)|^2 + \gamma |v(x, t)|^2) dx \leq W(0) + M \left(\frac{CaM}{|D_L|} + K - \log M \right).$$

Because of $\|u_0\|_{L^1(\Omega)} = M < 8\pi/\alpha\chi$, we can take a constant a satisfying

$$\left(a - \frac{\chi}{2}\right) \frac{1}{\alpha} - \frac{Ma^2}{16\pi} > 0.$$

Then gives

$$\sup_{0 \leq t < T_{max}} \int_{D_L} (|\nabla v|^2 + \gamma |v|^2) dx < \infty.$$

and hence $T_{max} = \infty$ by the case (ii) of Proposition 1. \square

3 Proof of Theorem 2

Proposition 3 *Let $h > 0$ and $\mathcal{E} = \{w \in C^1(\overline{D_L}) | \frac{\partial w}{\partial n} = 0 \text{ on } \partial D_L \text{ and } \|w\|_{L^1} \leq h\}$. Then, for any $\mathcal{F} \subset \mathcal{E}$ satisfy 1 or 2:*

(i) *For any $\varepsilon > 0$, there exists a positive constant C_ε s.t.*

$$\log \left(\int_{\Omega} e^w dx \right) \leq \frac{1 + \varepsilon}{16\pi} \int_{\Omega} |\nabla w|^2 dx + C_\varepsilon \text{ for } w \in \mathcal{F}.$$

(ii) *There exist a sequence $\{w_k\} \subset \mathcal{F}$, a point $q \in \partial D_L$, a constant $m \in [1/2, 1)$ and a regular measure ψ s.t.*

$$w^* - \lim_{k \rightarrow \infty} \frac{\exp(w_k) p_*}{\int_{D_L} \exp(w_k) p_* dx} = m \delta_q + \psi \quad \text{in } \mathcal{M}(\overline{D}),$$

where $p_* = \frac{8}{L^2(1+(x/L)^2)^2}$.

Next we consider the following elliptic problem related to the second equation of (P).

$$(EE) \quad \begin{cases} 0 = \Delta w - \gamma w + f, & x \in \Omega, \\ \frac{\partial w}{\partial n} = 0, & x \in \partial \Omega \end{cases}$$

with $f \in L^1(\Omega), \geq 0$.

Proposition 4 *Let $q \in \Omega$. Then there exists a positive constant η_0 such that for $\eta \in (0, \eta_0)$ there exists a positive constant C_{η_0} satisfying*

$$\int_{\Omega \cap B(q, \eta)} e^w dx \leq \exp \left(\frac{C_{\eta_0}}{\eta_0 - \eta} \|f\|_{L^1(\Omega)} \right) \int_{|x| < 2\eta_0} \frac{dx}{|x|^\theta},$$

where

$$\theta = \begin{cases} \frac{1}{2\pi} \int_{B(q, \eta_0)} |f| dx & \text{if } q \in \Omega, \\ \frac{1 + O(\eta_0)}{\pi} \int_{\Omega \cap D(q, \eta_0)} |f| dx & \text{if } q \in \partial\Omega. \end{cases}$$

Proof of Theorem 2: Let a be a positive constant satisfying

$$\left(a - \frac{\chi}{2}\right) \frac{1}{\alpha} - \frac{Ma^2}{16\pi} > 0.$$

We assume that $\{av\}$ satisfies 1 of Proposition 3, by which together with the arguments of proof of Theorem 1 it follows that $T_{max} = \infty$. It is the contradiction. Then, for any positive constant a with

$$\left(a - \frac{\chi}{2}\right) \frac{1}{\alpha} - \frac{Ma^2}{16\pi} > 0,$$

we observe that

$$\begin{aligned} \infty &= \limsup_{t \rightarrow T_{max}} \int_{\Omega} e^{av} dx \\ &\leq \frac{1}{m} \limsup_{t \rightarrow T_{max}} \int_{\Omega \cap B(q, \varepsilon)} e^{av} dx \quad \text{for any } \varepsilon > 0, \end{aligned} \quad (2)$$

by which together with Proposition 4 we have this theorem. \square

4 Proof of Theorem 3

By using Proposition 1 and Lemmas 1 and 2, we have (2) for any $q \in \mathcal{B}_I$ and any $a > \chi/2$, by which together with Proposition 4 we have this theorem.

References

- [1] H. Brezis and W. A. Strauss, Semi-linear second-order elliptic equations in L^1 , J. Math. Soc. Japan **25** (1973), 565–590.
- [2] M. A. Herrero and J. J. L. Velázquez, Singularity patterns in a chemotaxis model, Math. Ann., **306** (1996), 583–623.
- [3] M. A. Herrero and J. J. L. Velázquez, Chemotaxis collapse for the Keller-Segel model, J. Math. Biol., to appear.
- [4] W. Jäger and S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, Trans. Amer. Math. Soc., **329** (1992), 819–824.

- [5] E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an instability, *J. Theor. Biol.*, **26** (1970), 399–415.
- [6] T. Nagai, Blow-up of radially symmetric solutions to a chemotaxis system, *Adv. Math. Sci. Appl.*, **5** (1995), 581–601.
- [7] T. Nagai, T. Senba and K. Yoshida, Application of the Trudinger-Moser Inequality to a Parabolic System of Chemotaxis, *F. A.*, to appear.
- [8] A. Yagi, Norm behavior of solutions to the parabolic system of chemotaxis, *Math. Japonica*, to appear.